

Lecture 5

5.1 Stokes' and Green's Theorem

Before developing some more tools required to prove the Cauchy-Goursat theorem, we give some more motivation towards the result; what happens for C^1 functions. Recall Stokes's theorem for \mathbb{R}^2 : Consider a simply connected region $\Omega \subseteq \mathbb{R}^2$ with piecewise smooth, simple and closed boundary $\partial\Omega$. If $f = P dx + Q dy$ is a C^1 1-form on an open set $U \supseteq \Omega \cup \partial\Omega$, then

$$\oint_{\partial\Omega} f = \iint_{\Omega} df.$$

Now, as $f = P dx + Q dy$, we get $df = dP dx + dQ dy = P_y dy dx + Q_x dx dy = (Q_x - P_y) dx dy$. Hence,

$$\oint_{\partial\Omega} f = \oint_{\partial\Omega} P dx + Q dy = \iint_{\Omega} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

which is nothing but the statement of Green's theorem. We now return to \mathbb{C} and assume $f \in \text{Hol}(\Omega) \cap C^1(\Omega)$, where Ω is a simply connected region as above. Let $\gamma \subseteq \Omega$ be a piecewise smooth, simple and closed curve. Then, by the discussion of lecture 3,

$$\oint_{\gamma} f dz = \left(\oint_{\gamma} u dx - v dy \right) + i \left(\oint_{\gamma} v dx + u dy \right)$$

Using Stokes's theorem on the two integrals on the right we get, [Σ_{γ} is the capping surface of γ , in some lecture we have used the notion of capping surface as $\text{int}(\gamma)$]

$$\oint_{\gamma} f dz = \iint_{\Sigma_{\gamma}} \left(-\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy + i \iint_{\Sigma_{\gamma}} \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) dx dy = 0$$

where the final equality follows from the Cauchy-Riemann equations. This is exactly the statement of the Cauchy-Goursat theorem! We now set to develop the tools needed to remove the C^1 restriction, that is, to show that holomorphic functions must be C^1 on "nice" domains.

Exercise: Suppose $f \in C^1(\Omega)$, where Ω is as above. If γ is a curve as above, show that

$$\oint_{\gamma} f dz = 2i \iint_{\Sigma_{\gamma}} \bar{\partial} f dx dy$$

holds for all such functions, without assuming holomorphicity.

Recall the result from lecture 3 that

$$\frac{1}{2\pi i} \oint_{C_r(0)} \frac{1}{z} = 1$$

for any $r > 0$. Changing variables, we get the equality

$$\frac{1}{2\pi i} \oint_{C_r(z_0)} \frac{1}{z - z_0} = 1$$

for any $z_0 \in \mathbb{C}$. We now generalise this result to the following very useful theorem.


5.2 Cauchy Integral Formula

Theorem 5.2.1 (Cauchy Integral Formula) Let $f \in \text{Hol}(\mathcal{O})$ and $C \subseteq \mathcal{O}$ be a circle such that $D = C \cup \Sigma C$ is in \mathcal{O} . Then, for all $z \in \Sigma C$,

$$\frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta = f(z).$$

Remark. Note that if $z \in \mathcal{O} \setminus D$, by Theorem 4.1,

$$\frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta = 0.$$



Proof. Fix $z \in \Sigma C$ and let $B_\varepsilon(z) \subseteq \Sigma C$, $C_\varepsilon = \partial B_\varepsilon(z)$. We claim,

$$\oint_C \frac{f(\zeta)}{\zeta - z} d\zeta = \oint_{C_\varepsilon} \frac{f(\zeta)}{\zeta - z} d\zeta$$

Introduce 4 “cuts” as shown by the dotted lines in the figure, and let the loops displayed be oriented anticlockwise. Then, $\gamma_1 \cup \gamma_2 \cup \gamma_3 \cup \gamma_4 = C \cup (-C_\varepsilon)$. Further,

$$\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$$

is holomorphic in $\Sigma\gamma_j$ for each j . As each of these sets $\Sigma\gamma_j$ can be covered by an open convex set $\Omega_j \subseteq \mathcal{O}$, we get by Theorem 4.1 that

$$\oint_{\gamma_j} \frac{f(\zeta)}{\zeta - z} d\zeta = 0$$

for each j . Therefore, as integration over $\cup\gamma_j$ is simply the sum of each of these integrals, we get

$$\begin{aligned} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta &= \oint_{C_\varepsilon} \frac{f(\zeta)}{\zeta - z} d\zeta \\ \implies \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta - f(z) &= \frac{1}{2\pi i} \oint_{C_\varepsilon} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{f(z)}{2\pi i} \oint_{C_\varepsilon} \frac{1}{\zeta - z} d\zeta \\ \implies \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta - f(z) &= \frac{1}{2\pi i} \oint_{C_\varepsilon} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta \end{aligned}$$

By the triangle inequality,

$$\left| \frac{1}{2\pi i} \oint_{C_\varepsilon} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta \right| \leq \frac{1}{2\pi} \sup_{\zeta \in C_\varepsilon} \left| \frac{f(\zeta) - f(z)}{\zeta - z} \right| \ell(C_\varepsilon) = \varepsilon \sup_{\zeta \in C_\varepsilon} \left| \frac{f(\zeta) - f(z)}{\zeta - z} \right|$$

As $\varepsilon \rightarrow 0$, the supremum goes to $|f'(z)|$ and so the right most quantity goes to 0. Therefore, in light of the equality above, we get

$$\frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta - f(z) = 0$$

as was to be shown. □