

# Lecture 7

In this lecture, we will complete our discussion of power series and consider analytic functions over the complex plane. We will then prove the equivalence of analytic and holomorphic functions - deriving the Cauchy integral formulae in the process. This is truly marvelous result with no immediate analogue in real analysis, and will be extremely useful in the discussions to follow.

## 7.1 Power Series continued

We started our discussion of power series over  $\mathbb{C}$  in the previous lecture. We give some simple illustrations before moving on.

### Example 7.1.1

Consider the power series given by  $\sum_{n=1}^{\infty} (1 + (-1)^n) z^n$ . Then the radius of convergence for the same is given by

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} \sqrt[n]{|1 + (-1)^n|} = 1$$

Note that  $\lim \left| \frac{a_{n+1}}{a_n} \right|$  does not exist here, so one cannot use the ratio test.

### Example 7.1.2 (Exponential function)

Consider the power series  $\sum_{n=1}^{\infty} \frac{z^n}{n!}$ . Evidently,  $\lim \left| \frac{a_{n+1}}{a_n} \right| = 0$ , and thus the power series defines a function over  $\mathbb{C}$ . We call this the exponential function and denote it by  $\exp\{z\}$ . Also for  $z_1, z_2 \in \mathbb{C}$ , we have

$$\begin{aligned} \exp\{z_1\} \exp\{z_2\} &= \left( \sum_{k=0}^{\infty} \frac{z_1^k}{k!} \right) \left( \sum_{l=0}^{\infty} \frac{z_2^l}{l!} \right) \\ &= \sum_{n=0}^{\infty} \sum_{r=0}^n \frac{z_1^r z_2^{n-r}}{r!(n-r)!} \\ &= \sum_{n=0}^{\infty} \frac{(z_1 + z_2)^n}{n!} = \exp\{z_1 + z_2\} \end{aligned}$$

Note that the reordering of sums is justified as the series is absolutely convergent. Using this multiplicative property, along with continuity of power series, one can show that

$$\exp\{z\} = e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!},$$

where  $e := \exp\{1\} = \sum_{n=0}^{\infty} \frac{1}{n!}$ . We will talk about this in more detail in the upcoming lectures.

**Example 7.1.3** (Sine and Cosine functions)

We define the sine and cosine functions over  $\mathbb{C}$  using the complex power series analogous to the Taylor series expansion of their real counterpart. In particular,

$$\begin{aligned}\sin(z) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1}, \text{ and} \\ \cos(z) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} z^{2n}\end{aligned}$$

Note that these power series absolutely converge over  $\mathbb{C}$ , and thus define entire functions. We also have the Euler's identity

$$\begin{aligned}e^z &= \sum_{n=0}^{\infty} \frac{z^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} z^{2n} + i \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1} \\ &= \cos(z) + i \sin(z)\end{aligned}$$

We now comment on the differentiability of functions defined by power series. The exactly analogous result holds for real power series as well, and the proof is almost identical.

**Theorem 7.1.1**

Let  $R$  be the radius of convergence of the power series  $f(z) := \sum a_n z^n$ , where  $z \in B_R(0)$ . Then  $f \in \mathbf{Hol}(\mathcal{O})$  and the derivative  $f'(z) = \sum n a_n z^{n-1}$  is given by the corresponding term-by-term differentiation, for  $z \in B_R(0)$ .

*Proof.* First we compute the radius of convergence  $R'$  of the derived power series  $\sum n a_n z^{n-1}$ . Evidently,

$$\frac{1}{R'} = \limsup_{n \rightarrow \infty} \sqrt[n]{n a_n} = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n} = \frac{1}{R}$$

Thus the derived series defines a function  $g : B_R(0) \rightarrow \mathbb{C}$ , such that

$$g(z) = \sum_{n=0}^{\infty} n a_n z^{n-1}$$

Fix  $z \in B_R(0)$  and correspondingly choose  $\delta > 0$  such that  $|z| + \delta < R$ . Then for  $|h| < \delta$ ,

$$\frac{f(z+h) - f(z)}{h} - g(z) = h \sum_{n=2}^{\infty} a_n p_n(z, h),$$

where  $p_n(z, h) = \sum_{k=2}^n \binom{n}{k} h^{k-2} z^{n-k}$ . The proof now follows as

$$\begin{aligned}\left| \frac{f(z+h) - f(z)}{h} - g(z) \right| &\leq |h| \sum_{n=2}^{\infty} |a_n| |p_n(z, h)| \leq |h| \sum_{n=2}^{\infty} |a_n| p_n(|z|, \delta) \\ &\leq \frac{|h|}{\delta^2} \sum_{n=2}^{\infty} |a_n| (|z| + \delta)^n \rightarrow 0 \text{ as } h \rightarrow 0\end{aligned}$$

□

Thus, if  $R$  be the radius of convergence of a power series  $f(z) := \sum a_n z^n$ , where  $z \in B_R(0)$ , then  $f \in C^\infty(B_R(0))$ , where the power series coefficients are given by the corresponding term-by-term differentiation. Also, the coefficients  $\{a_k\}$  can be computed from the derivatives of the function  $f$  at  $z = 0$  as

$$f^{(k)}(z) = \sum_{n=k}^{\infty} n(n-1)\dots(n-k+1)a_n z^{n-k}$$

$$\therefore a_k = \frac{f^{(k)}(0)}{k!}$$

## 7.2 Analytic Functions

Polynomials constitute the simplest examples of functions holomorphic on a given domain. The next, most prototypical examples are those which are locally generated by power series.

### Definition 7.2.1 ► Analytic function

Let  $f : \mathcal{O} \rightarrow \mathbb{C}$  be a continuous function, where  $\mathcal{O}$  is open in  $\mathbb{C}$ . We say that  $f$  is analytic at  $z_0 \in \mathcal{O}$  if there exists  $r > 0$  such that  $f$  can be expressed as a convergent power series on  $B_r(z_0)$ . If this holds for all points  $z_0 \in \mathcal{O}$ , then we say  $f$  is analytic on  $\mathcal{O}$ .

From the results of the previous section, it is clear that analytic functions are holomorphic, and in fact, infinitely differentiable. For  $f : \mathcal{O} \rightarrow \mathbb{C}$  analytic at  $z_0$ , the local power series representation is

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k$$

The converse is decidedly false in the real case. Consider for instance, the function  $f \in C^\infty(\mathbb{R})$ , defined by

$$f(x) = \begin{cases} e^{-\frac{1}{x^2}} & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$$

It is easy to check that all derivatives of  $f$  vanish at  $x = 0$ , but  $f$  is not identically zero in any neighbourhood of  $x = 0$ . Thus,  $f$  is not analytic at  $x = 0$ . However this result has a positive answer for holomorphic functions; it truly is one of the most remarkable results of complex analysis.

**Theorem 7.2.1** Let  $\mathcal{O}$  be an open subset of  $\mathbb{C}$ , and  $f \in \mathbf{Hol} \mathcal{O}$ . Consider  $z_0 \in \mathcal{O}$  and  $\delta > 0$  such that  $\overline{B_\delta(z_0)} \in \mathcal{O}$ . Then, for  $C = C_r(z_0)$ , we have  $f(z) = \sum a_n (z - z_0)^n$ , where

$$a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$$

*Proof.* By Cauchy integral formula for a circular contour,  $\forall z \in B_r(z_0)$ ,

$$f(z) = \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta$$

Now, the following series converges uniformly for  $\zeta \in C \setminus \{z_0\}$ .

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - z_0} \frac{1}{1 - \left(\frac{z - z_0}{\zeta - z_0}\right)}$$

$$= \frac{1}{\zeta - z_0} \sum_{n=0}^{\infty} \left( \frac{z - z_0}{\zeta - z_0} \right)^n$$

Then, as Riemann integration behaves well under uniform convergence over compact domains, we have

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z_0} \sum_{n=0}^{\infty} \left( \frac{z - z_0}{\zeta - z_0} \right)^n d\zeta \\ &= \frac{1}{2\pi i} \sum_{n=0}^{\infty} (z - z_0)^n \oint_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \\ &= \sum_{n=0}^{\infty} a_n (z - z_0)^n \end{aligned}$$

Thus  $f$  is analytic over  $\mathcal{O}$ , with

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \oint_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \quad (7.1)$$

□

The equations (7.1) are known as Cauchy integral formulae. This theorem sets apart complex analysis from our familiar terrain of real analysis - and has many striking consequences. We will see many applications of this wonderful theorem in the next lecture. For now, we have the simple corollary:

#### Corollary

For  $\mathcal{O}$  an open subset of  $\mathbb{C}$ ,  $f \in \mathbf{Hol}(\mathcal{O})$  if and only if  $f$  is analytic on  $\mathcal{O}$ .